

Physics of Dense and Superdense Matter*

Mannque Rho^{a,b}

(a) Service de Physique Théorique, CE Saclay, 91191 Gif-sur-Yvette, France

(b) School of Physics, Korea Institute for Advanced Study, Seoul 133-791, Korea

(February 8, 2008)

Abstract

I discuss a few aspects of dense hadronic matter and superdense QCD matter that are considered to be relevant to the physics of compact astrophysical systems. The connection between a “bottom-up approach” and a “top-down approach” is made with the help of an effective field theory strategy. Topics treated are meson condensation going up from low density and color superconductivity going down from asymptotic density with the approach to the chiral phase transition made in terms of Brown-Rho scaling.

* Invited talk given at the First KIAS International Workshop on Astrophysics: Explosive Phenomena in Compact Astrophysical Systems, KIAS, Seoul, Korea; 24-27 May 2000.

1 Introduction

Super-dense hadronic and/or quark matter is considered to be relevant for understanding compact stars in the context of supernova explosions, hypernovae, merging of compact (neutron and black-hole) stars, gamma-ray bursts etc. See [1] for a recent review. It is not known at present precisely which densities are relevant to such systems but the region of the density involved is presumably somewhere in-between very low and very high density regimes, the two extreme regimes being in principle fairly well controlled. To study the physics of the density regime involved which cannot be systematically accessed as I will describe below, we have at our disposal basically two possible theoretical approaches: One, “bottom-up” one going up from very low density and the other, “top-down” going down from very high (asymptotic) density. The former relies on laboratory (experimental) data available at low density in nuclear and hadronic physics and the latter on the theoretical machinery available at super-high density at which QCD becomes weak-coupling. We would eventually like to bring the two approaches together and have them match. But both going up beyond nuclear matter density and going down from the asymptotic density to the relevant regime are difficult because of lack of reliable theoretical tools and of experimental data. In this talk I will discuss some aspects of both and indicate how the two “extremes” could be brought together at a density regime supposedly relevant to the astrophysical processes of interest. This talk will be qualitative, intended to give a personal overview of the present situation. Certain aspects of the topic somewhat lying outside of the main theme of this meeting are discussed in a separate note [2].

2 Phase Structure

In many of the current meetings in nuclear and hadronic physics, one is shown a canonical diagram depicting the phase structure of hadronic matter in terms of temperature (T) vs. chemical potential μ or baryon number density ρ . In heavy-ion meetings, the principal focus is put on the phase structure at high temperature. There, the phase below $T_c \sim 150$ MeV is in hadronic state and the one above in quark-gluon plasma. There is strong evidence from lattice gauge calculations that the phase transition does indeed take place at a given T_c although above the T_c , things are not as simple as the naive form of quark-gluon plasma that people have thought of before. In any event, it seems likely that one will be able to follow temperature properties of the systems with small non-zero density both theoretically and empirically, the former on lattice and the latter in the heavy-ion machines that are operating and/or are to come in the near future [3, 4]. The most recent phase diagram taken from Braun-Munzinger’s article [5] – which was also seen in Schukraft’s talk – is shown in Fig.1 [5]. It gives a perspective in this matter. There are many laboratories involved in this endeavor, such as “early universe,” LHC, RHIC, ..., SIS as indicated in the figure.

It is a different story with matter at high density. For technical reasons, it turns out to be extremely difficult to put dense matter on the lattice except for the academic case of $N_c = 2$ systems, accounting for the paucity of theoretical guidance as to what happens to hadronic systems at non-zero density. The knowledge we have up to date is based on various models. Models indicate that as density increases toward the point at which QCD (chiral) phase transition is supposed to take place, there can be a multitude of phase transitions that may or may not be in concordance with chiral phase transition. As I shall discuss below, there are some experimental data around nuclear matter density but there are no terrestrial data much

beyond. Thus there is need to resort to astrophysical observations. Neutron stars are the only laboratory available for this purpose as indicated in Fig.1. The aim of this note is to discuss how to proceed along the density axis of the figure.




Figure 1: The conjectured QCD phase diagram. It shows the recent development including the hadro-chemical freeze-out points determined from thermal model analyses of heavy-ion collision data at SIS, AGS and SPS energy. The hatched region represents the plausible phase boundary delineating two different phases of QCD indicated by lattice QCD calculations at $\mu_b=0$.

3 Probing Dense Matter

Matter at high temperature which may be relevant for the early Universe can be – and is being – probed in heavy-ion collisions at relativistic energy. Furthermore QCD at high temperature can be simulated on lattice which will eventually give a solid theoretical information. Now what about probing matter at high density? Up to nuclear matter density, ample information is available from nuclear systems. Thus combined with reasonable models, one could attempt to go beyond nuclear matter by extrapolating what we learn at nuclear matter density. This constitutes the “bottom-up” approach. At low density, hadrons are the proper variables in QCD and the “bottom-up” approach consists of doing systematic calculations in the strong-coupling regime aided by experimental data. However one cannot push this strategy too far: this scheme must break down at some density.

At asymptotic density, the theory is in principle well-defined since QCD is weak-coupling there and perturbation theory is well-defined. One may therefore start with the weak-coupling QCD and adopt the “top-down” approach. As we go down in density, however, the coupling

increases, so at some point the perturbative approach must also break down. The ultimate goal would then be to have the two approaches come together at some intermediate density. The difficulty is that the gap between the two break-down points is not narrow and it is in this gap that interesting astrophysical phenomena seem to take place. Furthermore, the region in question cannot be probed in terrestrial laboratory experiments. Thus no reliable predictions can be made. We believe that the compact stars such as neutron stars are the only systems from which we can gain empirical information for the regime concerned and since no lattice QCD is available at finite density at least in the near future, the compact stars seem to be the only direct sources for dense matter physics.

4 Bottom-Up Approach

4.1 Pions Do Not Condense

Beyond the normal matter density, the first phase change that is expected to occur is meson condensation. The lightest meson is the pion so one would think that pions would condense first [6]. What happens can be understood in a simple and caricatured way as follows.

Pions may condense in S or P waves. The S-wave condensation can occur due to a change in the pion mass in medium. This is a simple reflection of the fact that the mass term in scalar field theory is a “relevant” term. Since the pion mass in nature is ~ 140 MeV which is small compared with the chiral symmetry scale, $\Lambda_\chi \sim 1$ GeV, a small attraction in the scalar direction could push the mass down to zero, so one would think that it would be easy to trigger the phase transition with the order parameter $\langle \pi^a \rangle \neq 0$. Baryon matter density $\rho \sim \langle B^\dagger B \rangle$ where B is the baryon field could induce such a change which will be an S wave. In nature this phase transition does not seem to occur mainly because there is chiral symmetry protection. Although the pion is massive, its mass is small because chiral symmetry is broken explicitly by a small amount, i.e., the tiny quark mass which arises from the symmetry breaking at the electro-weak scale. Thus in some sense chiral symmetry is still operative and hence protects the pions from undergoing a phase change in the S-wave channel.

On the other hand, the P-wave condensation can occur due to the P-wave attraction that pions feel in the presence of nucleons. The interaction dictated by chiral symmetry is of the derivative type $\sim \frac{g_A}{f_\pi} \bar{N}^\dagger(x) \tau^a \sigma N(x) \cdot \nabla \pi^a(x)$ where g_A is the axial coupling constant and f_π is the pion decay constant, both of which are associated with the weak interaction. This P-wave interaction with the nucleon is strong as evidenced by the huge resonance $\Delta(1230)$ associated with the strong πN interaction. The attraction is greater the greater the coupling g_A . The P-wave attraction which increases with density in nuclear medium can generate instability against P-wave pion condensation signaled by a space-dependent order parameter $\langle \pi^a(x) \rangle$. What happens as density increases can be analyzed with a Lagrangian that preserves chiral symmetry structure of dense matter [7]. The crucial element in this formulation of pion condensation is that the phase transition is extremely sensitive to how g_A – which is associated with the space component of the axial current – behaves in nuclear medium. It turns out that the critical density goes like $\rho_c^\pi \propto (g_A^* - 1)^{-1}$. One can therefore get a qualitative idea as to how the P-wave condensation occurs by looking at how the effective coupling g_A^* behaves in nuclear medium.

There has been much controversy on the effective g_A in nuclei and nuclear matter and its relevance to chiral phase structure. Experiments in nuclei exciting giant Gamow-Teller resonances have been interpreted in terms of this effective Gamow-Teller coupling. Whatever may

be the valid interpretation, the outcome is that the *effective* g_A (that is, g_A^* which is density dependent)¹ tends to the limit

$$g_A^* \rightarrow 1 \quad \text{as } \rho \rightarrow \rho_0. \quad (1)$$

This argument does not pinpoint the critical density but one can say that ρ_c^π is pushed beyond the regime where the theoretical framework makes sense.

4.2 Kaon Condensation

As we will hear in the next talk by Chang-Hwan Lee, kaon condensation can occur in, and make an important impact on, neutron-star formation. I will give a very simplified argument for how and why kaons will condense in dense neutron-star matter. More rigorous discussion is given by Chang-Hwan Lee in [8].

In a nut-shell, a kaon, particularly the negative charged K^- which contains an anti-strange quark $\bar{s}$, can condense in neutron-star matter precisely because the strange quark is massive at the scale of low-energy physics we are concerned with. Since the strange quark has a current mass of ~ 150 MeV², chiral symmetry is explicitly broken in the strange sector and the low-energy theorems associated with Goldstone bosons sometimes fail badly. What the matter density does is to “rotate” the explicitly broken symmetry in such a way that the kaon mass gets lighter to partially restore the explicitly broken symmetry. Suppose that all Goldstone boson masses were zero. Then the same chiral symmetry protection argument that applies to pions will apply to kaons as well so that baryon number density cannot do anything to the energy density of the system. If on the other hand all Goldstone bosons were massive and degenerate, the rotation around the chiral circle (see Fig.2) would not affect the energy density and so nothing would happen. Indeed, things happen precisely because the potential is tilted due to the symmetry breaking and the baryon density does just the un-tilting to restore the symmetry triggering kaon condensation [10]. The transition is essentially triggered by the “relevant” mass term in the Lagrangian, so is S-wave. There is a beautiful support for this dropping-mass scenario from laboratory experiments [9].

In strong interactions, strangeness has to be conserved and if kaons were to condense, the pairs $K^\pm$ would have to do so. If it were to happen at a low enough density, say, $\rho/\rho_0 \sim 3$, then condensation could occur only if the repulsive term that counteracts the scalar attraction associated with the symmetry breaking term, namely, the so-called vector coupling, gets decoupled. In this case, both $K^\pm$ mesons become lighter and eventually may condense. But if this were to occur at much higher density than say $\sim 3\rho_0$, then kaons would be prevented from condensing in heavy-ion process for the reason to be mentioned below.

¹There has been much debate as to whether the quenching of g_A in nuclei and nuclear matter as evidenced in Gamow-Teller transitions is *fundamental* or some garden-variety nuclear phenomenon. I believe this is a non-issue. There is no question that taken as an *effective* parameter in a precisely defined sense, the g_A is quenched in nuclei. Whether it can be described by many-body effects (e.g., “core polarizations” due to tensor forces) or Δ -hole effects or even partial chiral restoration is mostly semantic – they are of the same physics – and is not relevant for the issue. What is relevant is that the *effective* Gamow-Teller matrix element can be described in dense medium by a quenched g_A^* in the chiral Lagrangian and that it goes to the value of 1 as density increases.

²The quark mass is not renormalization-group invariant and so has to be defined with a given scale. Even at a given scale, it is not well-determined. Various methods of analysis give the range 100 $\sim$ 200 MeV for the strange current-quark mass. This uncertainty makes the estimate of the critical density somewhat uncertain.




Figure 2: Projection onto the σ, K plane. The angular variable θ on the “chiral circle” represents fluctuation toward kaon mean field. Kaon condensation corresponds to “rotating” into the K direction (slightly) away from the chiral circle.

In the neutron-star matter, the kaon mass does not have to go all the way to zero. In fact when the electron chemical potential that increases with density – for non-relativistic nucleons – crosses the dropping kaon mass, the electron will be converted by weak interaction to the kaon

$$e^- \rightarrow K^- + \nu_e \quad (2)$$

and the condensing of kaons is favored for energetic reasons. Here K^+ will play no role. For K^- the vector interaction is attractive, so it speeds up the drop of the kaon mass.

The theoretical arguments that go into this treatment can be reliable only if the condensation occurs at low enough density. Were it to occur at much higher density, then there would be no reason to believe that the approximation makes any sense. Indeed the physics at higher density coming up in the bottom-up approach will be obstructed by the inability to handle the short-distance physics referred in nuclear physics to as short-range correlations. For instance a straightforward embedding of two-nucleon interactions at short distance into many-body systems would predict that the phase transition will take place at a much higher density than that relevant to neutron stars [11]. However this consideration is based on the Hartree approximation in a potential model with a KN potential that is not realistic at high density [12]. Nonetheless the issue of short-range correlations is open.

4.3 BR Scaling

Once kaons condense at some density ρ_{Kc} , going up further in density up to (QCD) chiral restoration is like opening a Pandora’s box. As mentioned, we have no firm proof for *any* transition with density since lattice calculations are not feasible at present. There are a large number of different and conflicting ideas published in the literature. I neither have time nor wish to go into details here. It is fair to say that nothing *additional* that is illuminating or startling is expected to emerge in the physics of compact stars in this regime, once the kaon condensation and related processes have done what Chang-Hwan describes in his talk. In this murky circumstance, I will take an option that allows me to go ahead all the same. That is to heed what is said in Chinese fortune cookies in North America: “*There are only two ways of doing things: One is the wrong way and the other is our way.*” I shall therefore take “our way” which is to invoke “Brown-Rho (BR) scaling” [13, 14].

Consider symmetric nuclear matter. (Asymmetric matter can be treated similarly.) At a density found in nuclear matter, one finds that a description in which all degrees of freedom

are *quasi-particles* treated in the mean field works well provided that those quasi-particles have masses and coupling constants that respect chiral symmetry and run with density [2, 15]. In particular if one considers an effective field theory with chiral symmetry with the nucleons, pions, scalar meson and vector mesons as the appropriate degrees of freedom, then with density, the masses run with density as ³

$$\Phi(\rho) \approx \frac{f_\pi^*(\rho)}{f_\pi} \approx \frac{m_\sigma^*(\rho)}{m_\sigma} \approx \frac{m_V^*(\rho)}{m_V} \approx \frac{M^*(\rho)}{M}. \quad (3)$$

Here M is the nucleon mass, and the subscripts σ and V stand respectively for the scalar σ and the vector mesons ρ, ω . Since the pion decay constant can be taken as an order parameter for chiral symmetry phases, the masses are all seen to follow the order parameter in density. Thus when $f_\pi^* \rightarrow 0$ in the chiral limit as $\rho \rightarrow \rho_\chi$ (where ρ_χ is the as yet undetermined critical density for chiral restoration), the masses approach zero in the same way. In nature, the chiral limit is not reachable, so the masses will not strictly go to zero but they will drop smoothly (in the scenario of BR scaling) as density increases with interesting physical consequences.

Following this BR scaling behavior all the way to the chiral transition clearly sidesteps the Pandora’s box alluded above. What happens in this scenario is then that as one approaches the chiral phase transition density, the relevant degrees of freedom will be quasi-quarks (rather than quasi-nucleons), pions, scalar σ (i.e., a chiral partner of the pion) etc. These will go *smoothly* over to the other side of the phase boundary. This smooth transition could also be witnessed when one comes down in the top-down way from asymptotic density as I will discuss below.

The BR scaling as postulated in [13] has support both from theory [2, 15] and from experiments, e.g., in inelastic electron scattering from nuclei and in dilepton production in heavy-ion collisions [12]. It seems to also account for the J/ψ suppression observed at CERN that is heralded to be the signal for the formation of quark-gluon plasma [16]. This may be one more evidence for “quark-hadron” continuity and/or “Cheshire Cat phenomenon” mentioned below.

5 Top-Down Approach

5.1 Color superconductivity

When the density is asymptotically high, QCD in weak coupling is applicable so one can compute things reliably. Because of asymptotic freedom, the gauge coupling becomes weak and the dynamics of the quarks is governed by one gluon exchange. In this situation, diquarks are mostly likely to condense in the scalar channel giving rise to color superconductivity. There have been many papers written recently on this subject [17]. I would like to discuss a few aspects of the phenomenon that may be relevant to the physics of compact stars.

When the quark chemical potential is so high that one can ignore the u-, d- and s-quark masses, the situation becomes simple and elegant. But as one goes down to the regime where the chiral phase transition takes place, the situation gets very muddy and we have no idea what’s really happening. Even color superconductivity scenarios differ depending upon how many flavors come into play. At non-asymptotic chemical potential, other excitations than diquark, such as quark-hole modes (e.g., Overhauser modes), can be equally or even more important. I

³These are not *pole* masses. They can however be related to physical masses given the appropriate corrections.

shall therefore focus on the ideal case of infinite chemical potential with three flavors of quarks and hope for the best as we go down to the density relevant to compact stars.

A left-handed quark is characterized by three colors (i.e., group $SU(3)_c$) and three flavors (i.e., group $SU(3)_L$), so can be denoted q_{La}^α with α standing for color and a for flavor and similarly for the right-handed quark. It turns out that it is most favorable to condense the diquark in the color-flavor-locked (CFL) state as

$$\langle q_{Li\alpha}^a(\vec{p}) q_{Lj\beta}^b(-\vec{p}) \rangle = - \langle q_{Ri\alpha}^a(\vec{p}) q_{Rj\beta}^b(-\vec{p}) \rangle = \kappa(p_F) \epsilon_{\alpha\beta} \epsilon^{abI} \epsilon_{ijI} \quad (4)$$

where i, j are the spin indices. This means that both color and chiral symmetries are spontaneously broken by the condensate as ⁴

$$SU(3)_c \times SU(3)_L \times SU(3)_R \rightarrow SU(3)_{c+L+R}. \quad (5)$$

As a consequence, seventeen Goldstone bosons are excited, i.e., nine scalars and eight pseudoscalars. Eight scalars get eaten up by the gluons which become massive and the remaining scalar gets condensed to give rise to superfluidity. The eight pseudoscalars constitute the lowest modes remaining in the system that pick up a mass proportional to the current quark mass that goes to zero as $\mu \rightarrow \infty$.

The most remarkable conjecture made by Schäfer and Wilczek [18] posits that there is one-to-one correspondence between the spectrum of excitations in the zero-density regime and that of excitations in the CFL phase. Apart from the 8 Goldstone pseudoscalars that are present in both regimes, the eight massive gluons possess the same quantum numbers as the octet vector mesons in free space and the 8 color-flavor-locked quarks (which may be simply understood as quark solitons or qualitons [19]) can be mapped to the octet baryons of zero density. The conjecture is that the infinite-density spectrum is *continuously connected* to the zero-density spectrum, implying “quark-hadron continuity” or in the more pictorial term, “Cheshire Cat phenomenon” [20]. This aspect shows that going toward high density beyond ρ_χ may reveal a richer and quite different variety than going to high temperature. There is however an obstacle to this intriguing possibility: there may be a series of phase changes as one goes down in density from infinite to the regime relevant to compact stars. First of all, even in color superconductivity (CSC), scenarios differ depending upon the scale of the strange quark mass relative to the chemical potential [17]. Not less importantly, there can be other phases than color superconductivity such as Overhauser effect [21], ISB [22] and others [23] that may intervene before going into the Goldstone phase of lower density. Despite the plethora of other possibilities, it is remarkable – and perhaps meaningful – that both pion condensation [24] and kaon condensation [25] could occur in the high density regime. If confirmed, this could be another support for the notion of quark-hadron continuity.

5.2 Astrophysical implications

The mere existence of pions and kaons in the CFL phase is a highly intriguing possibility one would like to explore. Although quarks/baryons are gapped in the CFL phase with a gap $\Delta \sim 10 - 100$ MeV, there can still be light fermions in some situations. These degrees of freedom could play an important role in astrophysical compact objects. Since it does not look feasible

⁴The $U(1)_V$ corresponding to the fermion number is also broken but the $U(1)_A$ symmetry is restored at high density. There is also a Z_3 invariance that remains but we need not be concerned with it here.

to make firm predictions for both terrestrial and astrophysical observables, one would like to turn the process around and *learn* from astrophysical observations about the state of matter in which such exotic phenomena take place.

There have been a series of papers addressing the role of color superconductivity (CSC) in the cooling of neutron stars [26]. The situation is quite complex and a complete analysis is probably not yet possible. The complexity comes from the fact that we do not really know what should be taken into account in the cooling since there are meson condensations and other processes that could intervene at the same time. Even within the framework of CSC, there can be several competing factors to consider. First there is the effect on the electro-weak interaction proper. A simple calculation shows that inelastic neutrino processes (i.e., neutrino production) can be enhanced in the CFL phase [27]. Next, the produced neutrinos may have longer mean-free path in the CFL phase than in normal phase [28], thereby increasing the emissivity. On the other hand, the gap Δ can be large so that there will be a strong suppression from the Boltzmann factor $\exp(-\Delta/T)$. All these competing factors would have to be carefully accounted for before one can come to any estimate of the cooling rate. A lot more work and help from observations will be needed for a clearer picture. We heard in this conference about a very interesting data coming from a point source in Cassiopeia A which could be a valuable clue to the issue [29].

Another interesting effect is on magnetic fields in compact stars [30, 31]. Since vector symmetry is spontaneously broken (actually in all scenarios of color superconductivity (CSC)), both color and ordinary electromagnetism are spontaneously broken. However although electrically charged diquarks condense as do electron pairs in ordinary superconductivity, they do not produce the familiar Meissner phenomenon of ordinary superconductivity. The reason is that there is an unbroken $U(1)$ symmetry with a massless gauge boson. What happens is that as long as the color gauge coupling g_c is much bigger than the electromagnetic coupling e , magnetic fields can penetrate into the system largely un-expelled from the CSC phase. As a consequence, if rotating compact stars are in the CSC phase, the magnetic field will be stable against decay on time scales longer than, say, 10^7 years set by observation [32]. In [30], it is even suggested that if an isolated pulsar were observed to lose its magnetic field as it spins down, it would rule out the presence of the CSC (or even quark) phase in the pulsar. Although quite interesting, this negative evidence is not too exciting. It would be a lot more exciting if the CFL or other CSC phases were associated with a positive consequence. Again, given the various other possibilities in the density regime involved, the situation may be different from this simplified consideration. It remains an open issue.

6 Summary

The approach to the density regime relevant to compact stars is made in both ways: “bottom-up” and “top-down.” The two extremes, namely, bottom and top, are fairly well described, the former based on effective field theory of QCD combined with experimental data and the latter based on weak-coupling theory of QCD. The two approaches leave a rather wide gap in between where they have no overlap and it is in this range that interesting phenomena seem to take place in dense matter in compact stars. A bridge is suggested in terms of BR scaling coming from bottom-up. Combined with what we learn from the top-down approach and from astrophysical observations, it would be possible to establish the intriguing “quark-hadron continuity” alias “Cheshire Cat.” The main aim of this talk was to stress that the compact stars

are the only source for physics at high density, the domain that cannot be accessed in terrestrial laboratories.

Acknowledgments

It is a pleasure to acknowledge the generous help and hospitality of Korea Institute for Advanced Study for my participation in co-organizing, and for attending, this first KIAS workshop on astrophysics and astro-hadron physics. This paper was prepared while I was visiting Korea Institute for Advanced Study.

References

- [1] Balberg, S., and Shapiro, S.L., astro-ph/0004317; Heiselberg, H., and Pandharipande, V., astro-ph/0003276.
- [2] Rho, M., nucl-th/0007060
- [3] See for a recent review with references, Satz, H., “Phase transitions in QCD,” hep-ph/0007209.
- [4] Schukraft, J., talk in this conference.
- [5] Braun-Munzinger, P., nucl-ex/0007021.
- [6] Migdal, A.B., Saperstein, E.E., Troitsky, M.A., and Voskresensky, D.N., Phys. Rept. **192**, 179 (1990).
- [7] See, e.g., Baym, G., and Campbell, D.K., “Chiral symmetry and pion condensation,” in *Pions in nuclei* ed. Rho, M., and Wilkinson, D.H., North-Holland, Amsterdam, 1978, pp. 1031-1094.
- [8] Lee, C.-H., Phys. Rep. **275**, 256 (1996)
- [9] Li, G.Q., Lee, C.-H., and Brown, G.E., Phys. Rev. Lett. **79**, 5214 (1997).
- [10] Brown, G.E., Kubodera, K., and Rho, M., Phys. Lett. **B175**, 57 (1987).
- [11] Carlson, J., Heiselberg, H., and Pandharipande, V., nucl-th/9912043.
- [12] Brown, G.E., and Rho, M., to appear in Phys. Rep.
- [13] Brown, G.E., and Rho, M., Phys. Rev. Lett. **66** (1991) 2720.
- [14] Brown, G.E., and Rho, M., Phys. Rep. **269**, 333 (1996).
- [15] Song, C., nucl-th/0006030, Phys. Rep., in press.
- [16] Tsushima, K., Sibirtsev, A., Saito, K., Thomas, A.W., and Lu, D.H., nucl-th/0005065; Phys. Lett. **B484**, 23 (2000).

- [17] For reviews, see Wilczek, F., “ QCD in Extreme Conditions,” hep-ph/0003183; Schäfer, T., “Color Superconductivity,” nucl-th/9911017; Rajagopal, K., “ Mapping the QCD Phase Diagram,” hep-ph/9908360, Nucl. Phys. **A661**, 150c (1999).
- [18] Schäfer, T., and Wilczek, F., Phys. Rev. Lett. **82**, 3956 (1999).
- [19] Hong, D.-K., Rho, M., and Zahed, I., Phys. Lett. **B468**, 261 (1999).
- [20] Nowak, M.A., Rho, M., and Zahed, I., *Chiral Nuclear Dynamics* (World Scientific, Singapore, 1996).
- [21] Park, B.-Y., Rho, M., Wirzba, A., and Zahed, I., Phys. Rev. **D62**, 034015 (2000).
- [22] Langfeld, K., and Rho, M., Nucl. Phys. **A660**, 475 (1999).
- [23] Kim, Y., and Rho, M., nucl-th/0005069.
- [24] Son, D.T., and Stephanov, M.A., hep-ph/0005225.
- [25] Schäfer, T., nucl-th/0007021.
- [26] Page, D., Prakash, M., Lattimer, J.M., and Steiner, A., hep-ph/0005094; Blaschke, D., Klähn, T., and Voskresensky, D.N., astro-ph/9908334.
- [27] Hong, D.-K., Lee, H.K., and Rho, M., to appear.
- [28] Carter, G.W. and Reddy, S., hep-ph/9995228.
- [29] Umeda, H., talk in this conference; Umeda, H., Nomoto, K., Tsuruta, S., and Nineshige, S., astro-ph/9910113
- [30] Alford, M., Berges, J., and Rajagopal, K., Nucl. Phys. **B571**, 269 (2000).
- [31] Blaschke, D., Sedrakian, D.M., and Shahabasyan, K.M., astro-ph/9904395.
- [32] Makashima, K., “Magnetic fields of binary X-ray pulsars,” in *The structure and evolution of neutron stars* eds. Pines, D. et al., Addison-Wesley, New York, 1992, p.86.